arXiv:1011.6025v1 [astro-ph.SR] 28 Nov 2010

Volume Title
*ASP Conference Series, Vol. **Volume Number***
Author
© ***Copyright Year** Astronomical Society of the Pacific*

Snooping around the big dog: VY CMa as seen with *Herschel*/HIFI

E. De Beck[1], L. Decin[1,2], K. M. Menten[3], A. Marston[4], D. Teyssier[4], and the HIFISTARS team

[1]*Department of Physics and Astronomy, Institute for Astronomy, K.U.Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium*

[2]*Astronomical Institute "Anton Pannekoek", University of Amsterdam, Science Park XH, Amsterdam, The Netherlands*

[3]*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*

[4]*European Space Astronomy Centre, ESA, P.O. Box 78, E-28691 Villanueva de la Cañada, Madrid, Spain*

Abstract. In the framework of the HIFISTARS guaranteed time key programme, we measured more than 70 molecular emission lines with high signal-to-noise ratio towards VY CMa using the high-resolution HIFI spectrometer (de Graauw et al. 2010) on board the *Herschel*[1] satellite. The kinematic information obtained from the measured water lines supports the hypothesis of multiple outflow components. The observed high-intensity maser lines give no indication for strong polarisation.

1. A simple intro to a complex outflow

VY CMa is a red supergiant at a distance of 1.1 kpc, with a luminosity of $3 \times 10^5\,L_\odot$, $T_{\rm eff} \sim 2800$ K, and a mass $\sim 25\,M_\odot$. Decin et al. (2006) derived a gas mass-loss rate of $\sim 3.2 \times 10^{-4}\,M_\odot$/yr. Millimeter interferometry reveals three kinematic components: *(1)* a dense, compact, and dusty central component, embedded in *(2)* a more diffuse and extended envelope, and *(3)* a high-velocity bipolar outflow (Muller et al. 2007).

2. Watery outflows

All H_2O lines in the left panel of Fig. 1, have very similar shapes and match a Gaussian profile centred at $v_{\rm LSR}$ = 21 km/s, with FWHM equal to the terminal velocity $v_{\rm e}$ derived from ground-based CO data, i.e. 46.5 km/s. Even higher velocities, up to 60 km/s, may be reached in the blue wing of the water lines (not shown in Fig. 1). Compared to a Gaussian we see strong self-absorption in the blue, excess emission in the red, and a "shoulder" around 10 km/s. The small bumps at the extreme blue velocities are the

---

[1] *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.



Figure 1. *Left* — comparing the shape of H_2O lines: *(a)* p-$H_2O(1_{1,1}$-$0_{0,0})$, *(b)* o-$H_2O(1_{1,0}$-$1_{0,1})$, *(c)* p-$H_2O(2_{1,1}$-$2_{0,2})$, *(d)* o-$H_2O(3_{0,3}$-$2_{1,2})$, *(e)* o-$H_2O(3_{1,2}$-$3_{0,3})$, *(f)* o-$H_2O(3_{1,2}$-$2_{2,1})$, *(g)* o-$H_2O(3_{2,1}$-$3_{1,2})$, *(h)* p-$H_2O(4_{2,2}$-$4_{1,3})$, *(i)* o-$H_2O(5_{3,2}$-$5_{2,3})$. The dashed profiles are Gaussian as described in the text. *Middle* — Gaussian-subtracted profiles. *Right* — selected maser lines: *(j)* SiO(15-14,v=1), *(k)* o-$H_2O(5_{3,2}$-$4_{4,1})$. The vertical dashed lines indicate the region $[\upsilon_{\rm LSR} - \upsilon_{\rm e}; \upsilon_{\rm LSR} + \upsilon_{\rm e}]$.

contributions from the outer layers moving toward the observer. This is as expected for the optically thick case. The thermal water emission thus likely traces all three outflow components.

3. Strong maser lines

Four strong maser lines were detected with HIFI: two rotational transitions of SiO in the first vibrationally excited state (13-12,v=1) and (15-14,v=1), and two of H_2O, both in the ground-vibrational state. Neufeld & Melnick (1991) predicted p-$H_2O(5_{2,4} - 4_{3,1})$ and o-$H_2O(5_{3,2} - 4_{4,1})$ to be maser lines. Their respective observed peak strengths are 3.4 K and 1.8 K in main beam temperature. It is remarkable that the o-H_2O maser was observed twice, with an interval of three weeks, and no variation was detected on this timescale (Harwit et al. 2010). Apart from the Goldreich-Kylafis effect in the wings of the o-H_2O maser, mentioned by Harwit et al. (2010), we did not detect any strong polarisation effects for any of these maser lines.

Acknowledgments. E. De Beck and L. Decin wish to acknowledge the financial support by FWO under grant number G.0470.07.

References

de Graauw, T., Helmich, F. P., Phillips, T. G., & others 2010, A&A, 518, L6+
Decin, L., Hony, S., de Koter, A., et al. 2006, A&A, 456, 549. `arXiv:astro-ph/0606299`
Harwit, M., Houde, M., Sonnentrucker, P., et al. 2010, A&A, 521, L51+. `1007.0905`
Muller, S., Dinh-V-Trung, Lim, J., Hirano, N., Muthu, C., & Kwok, S. 2007, ApJ, 656, 1109
Neufeld, D. A., & Melnick, G. J. 1991, ApJ, 368, 215